Pain Assessment and Documentation Tool

Introduction

The Pain Assessment and Documentation Tool (PADT) is a two-sided chart note designed to be easily included in a patient's medical record and to facilitate ongoing evaluation of patient pain and documentation of pain management. The PADT is intended to be administered by a clinician and includes sections to assess pain-related outcomes in four areas: analgesia, activities of daily living, adverse events (i.e., side effects), and aberrant drug-related behavior.

Progress Note Pain Assessment and Documentation Tool (PADT™)										
Patient Name: Reco	Patient Stamp Here									
Assessment Date:										
Current Analgesic Regimen										
Drug Name Strength (eg, mg)	Frequency	Maximum Total Daily Dose	:							
			_							
The PADT is a clinician-directed interview; that is, the clinician asks the questions, and the clinician records the responses. The Analgesia, Activities of Daily Living, and Adverse Events sections may be completed by the physician, nurse practitioner, physician assistant, or nurse. The Potential Aberrant Drug-Related Behavior and Assessment sections must be completed by the physician assistant , or nurse. The Potential Aberrant Drug-Related Behavior and Assessment sections must be completed by the physician . Ask the patient the questions below, except as noted.										
Analgesia	Activities of	 Daily Living	_							
If zero indicates "no pain" and ten indicates "pain as bad	Please indicate whether the pa	· · ·	_							
 as it can be," on a scale of 0 to 10, what is your level of pain for the following questions? 1. What was your pain level on average during the past week? (Please circle the appropriate number) 	current pain reliever(s) is Better, the Same, or Worse since the patient's last assessment with the PADT.* (Please check the box for Better, Same, or Worse for each item below.)									
No Pain 0 1 2 3 4 5 6 7 8 9 10 Pain as bad as it can be	Physical functioning	Better Same Worse								
What was your pain level at its worst during the past week?No Pain 0 1 2 3 4 5 6 7 8 9 10 Pain as bad	2. Family relationships									
as it can be 3. What percentage of your pain has been relieved	3. Social relationships									
during the past week? (Write in a percentage between 0% and 100%.)	4. Mood									
4. Is the amount of pain relief you are now obtaining from your current pain reliever(s) enough to make a real difference in your life?	5. Sleep patterns									
☐Yes ☐No	6. Overall functioning									
5. Query to clinician: Is the patient's pain relief clinically significant? Yes	*If the patient is receiving his or her first PADT assessment, the clinician should compare the patient's functional status with other reports from the last office visit.									

Copyright Janssen Pharmaceutica Products, L.P. ©2003 All rights reserved.

Progress Note Pain Assessment and Documentation Tool (PADT™)

1. Is patient experiencing any side effects from current discon						Potential Aberrant Drug-Related Behavior This section must be completed by the physician Please check any of the following items that you discovered during your interactions with the patient. Please note that some of these are directly observable			
				_		(eg, appears intoxicated), while others may require more active listening and/or probing. Use the "Assessment"			
						section below to note additional details.			
		None	Mild	Moderate	Severe	Purposeful over-sedation			
a. N	Nausea					Negative mood change			
						Appears intoxicated			
b. \	omiting/					Increasingly unkempt or impaired			
						Involvement in car or other accident			
c. (Constipation					Requests frequent early renewals			
						Increased dose without authorization			
d. I	tching					Reports lost or stolen prescriptions			
						Attempts to obtain prescriptions from other doctors			
e. N	Mental cloudiness					Changes route of administration			
						Uses pain medication in response to situational stressor			
f. S	Sweating					Insists on certain medications by name			
						Contact with street drug culture			
g. F	atigue					Abusing alcohol or illicit drugs			
						Hoarding (ie, stockpiling) of medication			
h. [Prowsiness					Arrested by police			
						☐ Victim of abuse			
i. (Other					Other:			
j. (Other								
2. P	atient's overall seve	erity of s	ide effe	cts?					
	None Mild	Mod		Severe					
	essment: (This secti			•	· ·	•			
-	our overall impressi oid therapy?	Yes	הווים ממוו	No		benefits, such as pain relief, outweigh side effects) from sure			
Cor	nments:								
Specific Analgesic Plan: Comments:									
Continue present regimen							•		
Adjust dose of present analgesic									
Switch analgesics Add/Adjust concomitant therapy									
Discontinue/taper off opioid therapy									
Date: Physician Signature:									